

Upper Aparima Catchment Group

Fresh water health and landscape influences
in Upper Aparima Catchment



**THRIVING
SOUTHLAND**

*Tonui ana te whenua. Tonui ana te takata.
A thriving, prosperous land. A thriving, prosperous people.*

AS AT OCTOBER 2021
Further updates will be included as new
information becomes available.

Welcome to Upper Aparima Catchment Brochure

This Upper Aparima Catchment Group brochure is one of a series of brochures on catchments in Southland.

Catchment Groups have been asking for more detailed insights into their individual catchments. This brochure provides insights based on available information, bringing together published science, research, data and information on the state of water, soil and land in the Upper Aparima Catchment.

It provides details on what affects water quality and how these impact the rest of the catchment, including out-of-catchment areas that may be impacted by what goes on in the catchment, such as the estuary.

Although this brochure collates all the available information that has been brought together in a literature review commissioned by Thriving Southland - called the Science Report thrivingsouthland.co.nz/science-report - it may not have all the details you know about in your catchment or the research you may have done on your farm or in your Catchment Group.

How to use this brochure

This brochure sets out publicly available details on:

- » Water quality
- » Landscape influences
- » Physiographic zones
- » Groundwater management zones (GMZs)
- » Measuring what lives in streams and rivers
- » Macroinvertebrate community index (MCI)
- » Estuary health
- » Where to get more information.

If you are not familiar with the terms and language used, read the brochure in conjunction with this glossary environment.govt.nz/publications/environment-aotearoa-2019-glossary.

We also recommend you check out the Catchment Group page on thrivingsouthland.co.nz/upper-aparima to learn more about the catchment and what projects the Catchment Group has underway or planned.

Interpreting what the data in the brochure means

Because this brochure brings together the data available, we have deliberately not interpreted that data or explained what the trends may mean for your catchment.

We recommend you contact an environmental consultant, your Thriving Southland Catchment Group coordinator or Environment Southland to speak to experts who can explain what these trends and data may mean for your catchment, or for your farm specifically.

You can also check out a range of information on the Thriving Southland Information Resource Hub thrivingsouthland.co.nz/infohub which will connect you with tools and resources from many different organisations to help you with understanding limit setting, environmental contributing factors, mitigations and options available to you.

A little bit about Thriving Southland

Thriving Southland supports Southland's Catchment Groups to understand challenges and opportunities in their catchments and create innovative and exciting solutions.

We have a vision to create a prosperous Southland, healthy people, and a healthy environment, and believe that by working together, Thriving Southland's communities will create a better future for all by protecting the region's prosperity, heritage, environment and health.

Thanks

Thank you to the farmers who supported the development of this brochure, and to the Ministry of Primary Industry for its Sustainable Land Use Programme which support the work Thriving Southland is delivering for farmers and communities in Southland.



Water quality in Aparima Catchment

Aparima Catchment

Upper Aparima is part of the Aparima Catchment which outflows into the Jacobs River Estuary. Aparima River and Jacobs River Estuary are an important source of mahinga kai, particularly shellfish, mussels, paua, tuna and inanga.

Currently Jacobs River Estuary is considered to be in fair to poor condition.

Summary of Upper Aparima Catchment


The hydrology, health and functions of a stream/river or area of groundwater are directly linked to the characteristics of its catchment, including the landscape, soils and human activities.

- » This is a large catchment covering a mosaic of six different physiographic zones.
- » Large areas overlie the Upper Aparima GMZ which has high nitrate levels in some locations.
- » Due to the cumulative effects from practices in Aparima Catchment, Jacobs River Estuary is considered a moderate eutrophication risk with indicators ranging from fair to poor with an overall expert opinion that the estuary is considered degraded.
- » Neighbouring farms in different zones may have very different water quality outcomes with similar farm practices due to different nutrient movements and pathways.



What does this mean?

- » In August 2020, the Government announced changes to the National Policy Statement for Freshwater Management (NPS-FM), the National Environmental Standard for Freshwater (NES) and changes to the Resource Management Act. These outline changes in regulations relating to wetlands and rivers, intensive winter grazing, intensification, stockholding areas and stock exclusion.
- » More about Environment Southland's response to the Government's Essential Freshwater package is here es.govt.nz/environment/water/essential-freshwater-package



Environment Southland, in partnership with Te Ao Mārama Inc*, has indicated a proposed limits and targets plan change (LTPC) will be notified in 2023. This will establish nutrient limits and targets to improve the quality of groundwater and surface water. waterandland.es.govt.nz/about/values-and-objectives

- » Environment Southland and Te Ao Mārama Inc have established a community-based regional forum to consider and advise on limits, targets, and methods.
- » Plan changes will result in additional controls and rules in Southland that will be focused on reducing the loss of nutrients, specifically nitrogen and phosphorus, and reducing discharges of sediment and faecal micro-organisms, from land to groundwater and surface water.
- » In the Environment Southland Proposed Water and Land Plan there is a focus on good management practices (GMPs) and farm environmental management plans (FEPs). You can view GMP factsheets for each physiographic zone on the Environment Southland website es.govt.nz

* Te Ao Mārama Incorporated looks after mana whenua interests in resource management and other aspects related to local government in Southland. It is authorised to represent three Ngāi Tahu papatipu runanga in Murihiku/Southland. It is involved in the protection of the spiritual and cultural values of the region, including wahi tapu (sacred places), mahinga kai (gathering of food and resources) and other natural resources.

Upper Aparima water quality

Surface water quality is assessed by testing how much nitrogen, phosphorus and *E. coli* is present. LAWA summary results for this catchment are (lawa.org.nz):

Total oxidised nitrogen

Monitoring site	5-year median	5-year trend	10-year trend	15-year trend
Aparima River at Dunrobin	0.027 mg/L			
Hamilton Burn at Affleck Road	0.51 mg/L			

^ Total Oxidized Nitrogen (TON) is the sum of nitrate and nitrite. Nitrite is generally a very small fraction of the TON concentration in rivers, TON is taken to be equivalent to the nitrate concentration

* 2016-2020 LAWA median per NPS-FM 2020 using TON as surrogate for NO₃-N

Too much TON can contribute to excessive algal growth in waterways.

Ammoniacal nitrogen

Monitoring site	5-year median	State	5-year trend	10-year trend	15-year trend
Aparima River at Dunrobin	0.005 mg/L	A			
Hamilton Burn at Affleck Road	0.005 mg/L	A			

If ammoniacal nitrogen reaches very high concentrations it can become toxic under certain temperature and pH conditions.

Dissolved reactive phosphorus

Monitoring site	5-year median	State	5-year trend	10-year trend	15-year trend
Aparima River at Dunrobin	0.002 mg/L	A			
Hamilton Burn at Affleck Road	0.005 mg/L	A			


Dissolved reactive phosphorus concentrations are an indication of a waterbody's ability to support nuisance algal or plant growths (algal blooms).

Total phosphorus


Monitoring site	5-year median	5-year trend	10-year trend	15-year trend
Aparima River at Dunrobin	0.004 mg/L			
Hamilton Burn at Affleck Road	0.014 mg/L			

Too much phosphorus can encourage the growth of nuisance plants such as algal blooms.

KEY (STATE)



KEY (TREND)



E. coli

Monitoring site	5-year median	State	5-year trend	10-year trend	15-year trend
Aparima River at Dunrobin	50n/100mL	A			
Hamilton Burn at Affleck Road	110 n/100mL	D			

* 2016-2020 LAWA median graded as per NPS-FM 2020

KEY (STATE)

A

Very good
(infection risk is 1%)

B

Good
(infection risk is 2%)

C

Fair
(infection risk is 3%)


D

Poor
(infection risk is >3%)


E

Very Poor
(infection risk is >7%)


KEY (TREND)




Very likely improving




Likely improving




Indeterminate



Very Likely degrading



Likely Degrading



Not Assessed

Results from lawa.org.nz (October 2021)

MCI

Macroinvertebrates include the caddisflies, mayflies, stoneflies, worms and snails that live in rivers. They are an important food source for fish and birds and sensitive to the combination of nutrients, sediment and habitat. Due to this sensitivity, they are considered to be a good representation of overall water quality and ecosystem health. The different macroinvertebrates present can be identified and then converted to a score called the MCI.

A higher MCI score generally indicates a healthier stream. Generally, MCI scores range from >150 (very good water quality) to as low as 20 (very poor water quality).

The MCI scores for Upper Aparima Catchment are (LAWA October 2021):

MCI

Monitoring site	5-year median	State	10-year trend	15-year trend
Aparima River at Dunrobin	118.0	B		
Hamilton Burn at Affleck Road	96	C		

KEY (STATE)

A

Macroinvertebrate community indicative of pristine conditions with almost no organic pollution or nutrient enrichment.

B

Macroinvertebrate community indicative of mild organic pollution or nutrient enrichment. Largely composed of taxa sensitive to organic pollution/nutrient enrichment.

C

Macroinvertebrate community indicative of moderate organic pollution or nutrient enrichment. There is a mix of taxa sensitive and insensitive to organic pollution/nutrient enrichment.

D

Macroinvertebrate community indicative of severe organic pollution or nutrient enrichment. Communities are largely composed of taxa insensitive to inorganic pollution/nutrient enrichment.

National bottom line: MCI score 90

KEY (TREND)

Very likely improving

Likely improving

Indeterminate

Very Likely degrading

Likely Degrading

Not Assessed

Estuary health

Table: Estuary state information (provided by Environment Southland July 2021 es.govt.nz state and outcome map).

Although the estuary is not in the Upper Aparima Catchment, it is an important factor in understanding the impacts of water quality in Upper Aparima. Decisions made in Upper Aparima that affect water quality upstream, flow downstream and impact on water quality in the estuary.

The below assessment of estuary health based on the sediment quality gives a good indication of what is happening upstream across all the catchments that feed into the waterways supplying the estuary and therefore impact on the ecosystems and the cultural values of the area.


Estuary	Soft mud	Nutrients in sediment	Oxygen in sediment	Macroalgae cover	Seagrass loss	GEZ*
Waiau Lagoon/Te Wae						
Lake Brunton						
Waituna Lagoon/Waiparera (not assessed)						
New River Estuary						
Jacobs River Estuary						
Waikawa Estuary						
Haldane Estuary						
Freshwater Estuary						
Waimatuku Estuary						
Toetoes Estuary						

KEY  Very Good  Good  Fair  Poor



* Gross Eutrophic Zone (GEZ) is defined as an area that has low sediment oxygenation (<1cm aRPD), soft mud (>25% mud content) and the presence of high macroalgal cover (>50% cover). These areas are in poor condition and can no longer support most estuarine animals and shellfish.

Aparima Catchment with Estuaries and Surface water quality monitoring sites*

Water quality state and MCI



Estuary risk indicators



*Sites in Upper Aparima Catchment only shown

Landscape influences

What we do on the land can affect our water, but how it affects our water depends on a range of factors, including how our landscape works. It is useful to look at:

- » Physiographic zones which help to explain how nitrogen, phosphorus, sediment and faecal micro-organisms (such as *E. coli*) move and are attenuated (reduced, e.g. by biological or chemical processes).
- » GMZs which highlight the connectivity between land, surface water and groundwater.

Physiographic zones

Southland has been divided into nine physiographic zones to help understand how water moves across the landscape and why water quality is better in some places than others. Each physiographic zone represents an area that has similar factors influencing water quality, such as climate, topography, geology and soil type.

Extensive areas of this catchment fall into the alpine and bedrock/hill country physiographic zones. Lower lying areas are mainly classified as oxidising, gleyed or peat wetlands. A minor component is the riverine physiographic zone, which extends along the margins of major rivers (see map). These zones differ in the way nutrients are transported and attenuated within the catchment.



Upper Aparima Catchment showing physiographic zones

Alpine

This zone occupies land above 800 metres elevation. It is steeply sloping with high rainfall and snow. There is limited loss of nutrients and faecal microbes from this zone due to low land use intensity.

Bedrock/hill country – overland flow

This zone is found on rolling to steep land below 800 metres. It is characterised by high rainfall and a dense network of branching streams.

Water quickly flows down-slope to nearby streams following high or prolonged rainfall. Nitrogen, phosphorus, sediment and faecal micro-organisms are all carried with water, particularly during late autumn and winter.

Riverine

Located along the margins of major rivers, this zone is characterised by shallow, stony soils that drain quickly to underlying shallow aquifers. This zone transports nutrients, particularly nitrogen, to coastal estuaries and lagoons.

Peat wetlands

This zone features poorly drained peaty soils that are extremely acidic. The water table in these areas is high. Developed areas require extensive artificial drainage. Soluble phosphorus concentrations are high in acidic oxygen depleted ground and surface waters; conversely nitrate concentrations are low.

Gleyed


This zone is generally found in areas that were once wetlands. It is characterised by a dense network of streams and a high water table during winter.

Soils are prone to waterlogging and have some denitrification* ability, which reduces build up of soil nitrogen. However, an extensive network of artificial drainage rapidly transports nitrogen, phosphorus, sediment and faecal microbes to surface water, particularly during heavy rain.

Oxidising

Soils and aquifers in this zone have high risk of nitrogen build-up due to low rates of denitrification*.

The combination of flat land and well drained soils results in high rates of nitrogen leaching (deep drainage) to underlying aquifers.



*Denitrification occurs when nitrate is converted to nitrogen gas via various reactions involving bacteria. Where denitrification occurs, nitrogen is effectively lost from soil and water as a gas. This is a form of attenuation.

GMZ - Upper Aparima

Large areas of Upper Aparima Catchment overlie the Upper Aparima GMZ.




Map Source: Environment Southland maps.es.govt.nz/index.aspx?app=water-and-land

Upper Aparima GMZ

The Upper Aparima GMZ covers an area of approximately 49,000ha in Aparima River Catchment upstream of Otautau:

- » Depth to groundwater in the Aparima GMZ typically ranges from less than two metres below ground level along the margins of Aparima River to more than 10 metres below ground level under elevated terraces toward the valley margins.
- » Seasonal variation in groundwater levels is generally less than two to three metres, reducing on lower terraces adjacent to the Aparima River.
- » See below for a diagrammatic cross-section of this GMZ showing areas of groundwater (source es.govt.nz/environment/water/groundwater/groundwater-management-zones/upper-aparima).
- » Groundwater recharge in this zone is derived from local rainfall and runoff from surrounding hills that soaks through the soil. Therefore there is a high risk of groundwater contamination from leaching.



Groundwater quality Upper Aparima GMZ

- » Nitrate = variable, with some areas having high concentrations
- » Phosphorus = low
- » *E. coli* = low, but risk may be elevated where soils are well-drained and the water table is shallow.



Find out more

Environment Southland Aparima flood warning fact sheet
bit.ly/3cLate5

Find out more about physiographic zones
bit.ly/2OI7z7F

Find out more about Southland's groundwater
bit.ly/30Db5g1

Find out more about stream health

Environment Southland
es.govt.nz/environment/water/rivers-and-streams

Land Air Water Aotearoa (LAWA)
lawa.org.nz

Ministry for the Environment
environment.govt.nz/facts-and-science/freshwater

Link to iwi freshwater objectives
bit.ly/2P4HsBV

Get in contact

For more information about your catchment and to contact your local catchment coordinator

021 466 700 | office@thrivingsouthland.co.nz
thrivingsouthland.co.nz/catchment-groups



THRIVING SOUTHLAND

Tōnui ana te whenua. Tōnui ana te takata.

A thriving, prosperous land. A thriving, prosperous people.